

# GENITOURINARY MEDICINE

Editor: A Mindel (Sydney)

President MSSVD: J S Bingham (London)

Hon Sec MSSVD: T McManus (London)

M W Adler (London)

R Ashley (Seattle)

D A Cooper (Sydney)

L Corey (Washington)

Y Cossart (Sydney)

B Donovan (Sydney)

J Douglas (Denver)

L M Drusin (New York)

J S Greenspan (San Francisco)

G Gross (Hamburg)

G R Kinghorn (Sheffield)

K Holmes (Seattle)

M Laga (Antwerp)

A S Latif (Harare)

D Mabey (London)

A Maheus (Antwerp)

Assistant Editor: C A Carne (Cambridge)

Assistant Editor, Abstracts: C Sonnex (Cambridge)

D Mercey (London)

A R Moss (San Francisco)

J Perriens (Geneva)

R S Pattman

(Newcastle-upon-Tyne)

A J Pinching (London)

A Pozniak (London)

G Ridgway (London)

J A R van den Hoek

(Amsterdam)

J Weber (London)

L Westrom (Lund)

Editor British Medical Journal

Technical Editor: Kenneth Till

(Taunton, UK) Fax (UK) 09847-315

Telephone (UK) 09847-388

This *Journal*, founded by the Medical Society for the Study of the Venereal Diseases, publishes original work on the investigation and treatment of genitourinary and allied disorders, and review articles, correspondence, and abstracts.

**ADVICE TO AUTHORS** Papers for publication, which will be accepted on the understanding that they have not been and will not be published elsewhere and are subject to editorial revision, should be sent in duplicate to **Prof A Mindel**, Academic Department of Sexual Health Medicine, Sydney Hospital, Macquarie Street, Sydney 2000, GPO Box 1614, Sydney, NSW, Australia. Tel: 61 2221 2800; Fax: 612 221 5810. All authors must give signed consent to publication. Submitted papers must be accompanied by the following statement, signed by all the authors: "No paper resembling the enclosed article has been or will be published except in *Genitourinary Medicine*. We transfer all copyright ownership to *Genitourinary Medicine*." If requested, authors shall produce the data upon which the manuscript is based for examination by the editor. Manuscripts will only be acknowledged if a stamped addressed postcard or international reply coupon is enclosed. Rejected papers will not be returned. Figures will be returned if requested at the time of submission.

Full details of requirements for manuscripts in the Vancouver style (*BMJ* 1982;284:1766-70) are given in *Uniform requirements for manuscripts submitted to biomedical journals*, available from the Publishing Manager, *British Medical Journal*, BMJ Publishing Group, BMA House (50p post free). Brief details are as follows:

(1) **SCRIPTS** (including correspondence and book reviews) must be typewritten on one side of the paper in double spacing with ample margins. Two copies should be sent.

(2) **EACH SCRIPT** should include, in the following order: a brief summary (structured summaries are preferred), typed on a separate sheet, outlining the main observations and conclusions followed by up to three key words suitable for use in indexing; the text divided into appropriate sections; acknowledgements; references; tables, each on a separate sheet; and legends for illustrations.

(3) **THE TITLE** of the paper should be as brief as possible.

(4) **THE NUMBER OF AUTHORS** should be kept to the minimum, and only their initials and family names used.

(5) **ONLY THE INSTITUTION(S)** where work was done by each author should be stated.

(6) **SI UNITS** must be used. If old fashioned units are used, SI units should be given in parentheses or, for tables and figures, a conversion factor given as a footnote.

(7) **ONLY RECOGNISED ABBREVIATIONS** should be used.

(8) **ACKNOWLEDGEMENTS** should be limited to workers whose courtesy or help extended beyond their paid work, and supporting organisations.

(9) **FIGURES** should be numbered in the order in which they are first mentioned in the text. Captions should be typed on a separate sheet. **DIAGRAMS**: use thick, white paper and insert lettering lightly in pencil. **PHOTOGRAPHS**: should be marked lightly on the back with the author's name and indicating the top, and should not be attached by paper clips or pins. They should be trimmed to include only the relevant section (and will be reproduced 68 or 145 mm wide) to eliminate the need for reduction. Photomicrographs must have internal scale markers. Radiographs should be submitted as photographic prints, carefully prepared so that they bring out the exact point to be illustrated.

(10) **TABLES** should be numbered, have titles, and be typed on separate sheets. Please avoid large tables and use

the format which appears in current issues.

(11) **REFERENCES** should be numbered consecutively the first time they are cited and identified by arabic numbers in the text, tables, and legends to figures. Authors must take full responsibility for the accuracy of their references, and the list should be kept as short as practicable. It should be in the order in which references are first mentioned, and should include (in the following order), *journals*: author's name and initials, title of paper, name of journal (in full or abbreviated according to the list in *Index Medicus*), year of publication, volume number, and first and last page numbers; *books*: author's name and initials, full title, edition, place of publication, publisher, and year of publication. When a chapter in a book is referred to, the name and initials of the author of the chapter, title of the chapter, "In:", name and initials of the editor, and "ed" should precede book title, etc as above. In references to journals or books, when there are seven or more authors the names of the first three should be given followed by "*et al.*" Names of journals no longer published or not in *Index Medicus* should be given in full—for example, *British Journal of Venereal Diseases*.

**PROOFS** Contributors receive one proof, and should read it carefully for printers' errors and check the tables, figures, legends, and any numerical, mathematical, or other scientific expressions. Alterations should be kept to a minimum.

**REPRINTS** A limited number of reprints may be ordered at cost from the Publishing Manager when the proofs are returned.

**NOTICE TO SUBSCRIBERS** This *Journal* is published bi-monthly. The annual subscription rates are £126.00 and \$237 in the USA. Subscribers may pay for their subscriptions by Access, Visa, or American Express by quoting on their orders the credit or charge card preferred, the appropriate personal account number, and the expiry date of the card. Orders should be sent to the Subscriptions Manager, *Genitourinary Medicine*, BMJ Publishing Group, BMA House, Tavistock Square, London WC1H 9JR. Orders can also be placed locally through any leading subscription agent or bookseller. (For the convenience of readers in the USA, subscription orders, with or without payment, can be sent to: *British Medical Journal*, Box 560B, Kennebunkport, Maine 04046. All inquiries, however, including those regarding air mail rates and single copies already published, should be addressed to the publisher in London.)

Second class postage paid, Rahway NJ Postmaster. Send address changes to: *Genitourinary Medicine* c/o Mercury Airfreight International Ltd Inc, 2323 Randolph Avenue, Avenel, NJ 07001, USA.

**NOTICE TO ADVERTISERS** Applications for advertisement space and for rates should be addressed to the Advertisement Manager, *Genitourinary Medicine*, BMJ Publishing Group, BMA House, Tavistock Square, London WC1H 9JR.

**COPYRIGHT** © 1994 by *Genitourinary Medicine*. This publication is copyright under the Berne Convention and the International Copyright Convention. All rights reserved. Apart from any relaxations permitted under national copyright laws, no part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior permission of the copyright owners. Permission is not, however, required to copy abstracts of papers or of articles on condition that a full reference to the source is shown. Multiple copying of the contents of the publication without permission is always illegal.

ISSN 0266-4348

## MATTERS ARISING

### Diagnosis of genital human papillomavirus lesions in the male

In an excellent study by Hippeläinen *et al*<sup>1</sup> the histopathological criteria for HPV infection were evaluated. As the authors point out HPV DNA may be detected in tissue lacking histopathological HPV associated signs and so can also the opposite situation occur.

Out of the HPV DNA negative biopsies (in situ hybridisation (ISH) technique) 26.7% did show koilocytosis, a sign said to be linked to a replicative HPV infection. The conclusion that koilocytosis is the strongest predictor for HPV positivity in flat genital lesions, giving a risk ratio of 3.7, is worthy of note but it would be interesting to hear the authors further discuss the finding of koilocytosis in the HPV DNA negative patients and the completely normal histology in the adjacent biopsied area. Were no pathological signs found at all? HPV are known to be latent in clinically normal tissue.<sup>2</sup>

As for the characteristic vacuolisation, also mentioned to be associated to HPV infection, maybe it should be discussed since it could be misinterpreted as koilocytosis by the pathologists not familiar with the signs of HPV infection. In the study referred to vacuolisation was more frequent in HPV negative cases than in HPV positive ones. This inverse correlation is only noticeable in table 4 but is for certain of interest. The authors conclude that HPV typing is essential for diagnosis in doubtful cases (lacking koilocytosis). We want to point out the fact that there is a diagnostic difficulty also in the case where koilocytosis is present and the rather insensitive ISH is negative.

ANDERS STRAND  
EVA RYLANDER  
University Hospital,  
Uppsala, Sweden

- 1 Hippeläinen MI, Syrjänen S, Hippeläinen MJ, Saarikoski S, Syrjänen K. Diagnosis of genital human papillomavirus (HPV) lesions in the male: correlation of peniscopy, histology and in situ hybridisation. *Genitourin Med* 1993;69:346-51.
- 2 Ferenczy A, Mitao M, Nagai N, Silverstein SJ, Crum CP. Latent papillomavirus and recurring genital warts. *N Engl J Med* 1985;313:784-8.

#### Hippeläinen *et al* reply:

While opening the discussion about our recent paper,<sup>1</sup> Drs Strand and Rylander focused on an important issue; the concordance between histology and HPV DNA typing in genital HPV lesions. Indeed, the sometimes low concordance between these two techniques as a source of diagnostic problems has been emphasised in the female genital lesions as well, but according to our experience, such a discordance seems to be an even more severe problem in the diagnosis of genital HPV infections in the males.

In the present series, a type-specific in situ hybridisation (ISH) with seven different

probes (HPV 6,11,16,18,31,33,42) was used as the detection method of HPV DNA.<sup>1</sup> As shown in table 4, the two morphological signs in the biopsies most frequently associated with HPV DNA-positivity were koilocytosis (67.6%) and acanthosis (61.7%). However, these two histological features were detected in HPV DNA-negative biopsies as well, that is, in 26.7% and 40.4% of cases, respectively. As pointed out by Strand and Rylander, koilocytosis is generally accepted as the cytopathic effect of HPV, being the hallmark of a productive HPV infection.

However, in diagnostic studies like this, it is frequently impossible to discover the specific HPV type inducing the koilocytotic change in all tissue biopsies, that is, to achieve a 100% correlation between morphology and presence of HPV DNA. This has multiple reasons. The two most feasible ones to explain the situation in the present series include the following: (1) Of the currently recognised 71 different HPV types, more than 30 are known to infect the genital mucosa. If all these probes are not included in the diagnostic test panel, it is more than probable that a minor or major proportion of the biopsy specimens analysed remain HPV DNA-negative. Our test panel included the seven most frequent HPV types, covering usually some 70% of the HPV-positive samples in most series.<sup>2</sup> (2) Albeit a highly applicable diagnostic tool, ISH has its limitations, namely, its sensitivity of approximately 20 viral copies/cell. Accordingly, any cell harbouring HPV DNA copies below that limit, most likely remains undetected by ISH. For some unknown reason, the viral load in the male genital HPV lesions seems to be substantially lower than that in the female HPV lesions, which might be one of the reasons why HPV DNA-detection rate by ISH is significantly higher in the latter.

The latter point (2) is also pertinent to the second issue raised by Strand and Rylander; the detection of HPV DNA in histologically normal epithelium. Of the 135 biopsies classified as normal on light microscopy, HPV DNA was found in nine (6.7%) (table 4). This is in agreement with the concept on latent HPV infections, the diagnostic criteria of which were recently outlined in detail.<sup>3</sup>

The third subject pointed out in their letter is the difficulty in making the distinction between true koilocytes and vacuolised cells. The former are considered as the cytopathic sign of HPV, the latter most probably arise as a result of multiple non-specific stimuli. In fact, such vacuolised cells were rare (6.9%) in HPV DNA-positive lesions, but more frequent (14.7%) in HPV DNA-negative biopsies. We completely agree with Strand and Rylander in their statement that the correct diagnosis of male genital HPV lesions is sometimes difficult both in cases with and without koilocytosis, even if you have ISH in use. It is good to remember, however, that even more difficult problems (albeit of different character) may arise, if more sensitive HPV DNA detection techniques, such as PCR are used.

MARITA HIPPELÄINEN  
STINA SYRJÄNEN  
MIKKO HIPPELÄINEN  
SEPPO SAARIKOSKI  
KARI SYRJÄNEN  
Kuopio University Hospital,  
Kuopio, Finland

- 1 Hippeläinen MI, Syrjänen S, Hippeläinen MJ, Saarikoski S, Syrjänen K. Diagnosis of genital human papillomavirus (HPV) lesions in the male: correlation of peniscopy, histology and in situ hybridization. *Genitourin Med* 1993;69:346-51.
- 2 Kataja V, Syrjänen K, Syrjänen S, *et al*. Prospective follow-up of genital HPV infections: Survival analysis of the HPV typing data. *Eur J Epidemiol* 1990;6:9-14.
- 3 Syrjänen KJ. Genital Human papillomavirus (HPV) infections and their associations with squamous cell cancer: reappraisal of the morphologic, epidemiologic and DNA data. In: Fenoglio-Preiser CM, Wolff M, Rilke F, eds. *Progress in Surgical Pathology*. Field & Wood 1992 Vol XII 217-240.

## NOTICES

### The UK Family Planning Association

The UK Family Planning Association is beginning a research project to explore the development of recent family planning and reproductive health initiatives within the UK. We particularly wish to gather information from people currently or recently associated with clinical facilities which combine access to family planning information and services with genitourinary medicine and sexual health provision. In the first instance we wish to invite health and social welfare professionals with experience of planning, developing, managing or working in combined clinic services to contact:

Joan Walsh  
Health Policy and Research Officer  
Family Planning Association  
27-35 Mortimer Street  
London W1N 7RJ, UK  
Tel: 071 636 7866

### Fifth Congress of the European Academy of Dermatology. 13-17 October 1996. Lisbon, Portugal

Further information may be obtained from: Professor A. Cabral Ascensão, Clínica Universitária de Dermatologia e Venereologia, Hospital Pulido Valente, Alameda das Linhas de Torres, 1700 Lisboa, Portugal.

### IUVDT World STD/AIDS Congress 1995. 19-23 March 1995, Singapore

For further information contact: Communication Consultants, 336 Smith Street #06-302, New Bridge Centre, Singapore 0105. Telephone (65) 227 9811. Fax (65) 227 0257.

## CURRENT PUBLICATIONS

Selected titles from recent reports published worldwide are arranged in the following sections:

Gonorrhoea  
Chlamydia  
Pelvic inflammatory disease  
Candidiasis  
Bacterial vaginosis  
Trichomoniasis  
Syphilis and other treponematoses  
Hepatitis  
Herpes  
Human papillomavirus infection  
Cervical cytology and colposcopy  
Other sexually transmitted diseases  
Public health and social aspects  
Microbiology and Immunology  
Dermatology  
Miscellaneous

### Gonorrhoea

**Multicenter trial of single dose azithromycin vs ceftriaxone in the treatment of uncomplicated gonorrhoea.**

HH HANDSFIELDS, ZA DALU, DH MARTIN, *et al.* *Sex Transm Dis* 1994;21:107.

**Bactericidal activity of six antibiotics against *Neisseria gonorrhoeae*.**

Y CRENN, M MEYRAN, T SAMSON, JD CAVALLO. *J Antimicrob Chemother* 1994;33:855.

**Variation in hydrogen peroxide sensitivity between different strains of *Neisseria gonorrhoeae* is dependent on factors in addition to catalase activity.**

TM ALCORN, HY ZHENG, MR GUNTHER, DJ HASSETT, MS COHEN. *Infect Immun* 1994;62:2138.

### Chlamydia

**Risk factors for recurrent *Chlamydia trachomatis* infections in women.**

SD HILLIS, A NAKASHIMA, PA MARCHBANKS, DG ADDISS, JP DAVIS. *Am J Obstet Gynecol* 1994;170:801.

**Chlamydia serologic studies and recurrent spontaneous abortion.**

R RAE, IW SMITH, WA LISTON, DC KILPATRICK. *Am J Obstet Gynecol* 1994;170:782.

**Genital chlamydial infection; diagnostic practices of general practitioners in Melbourne, Australia.**

F WESTGARTH, N CROFTS, DM GERTIG. *Sex Transm Dis* 1994;21:118.

**Factors associated with the geographic variation of reported chlamydia infection in Minnesota.**

EA BELONGIA, SJ MOORE, RS STEECE, KL MACDONALD. *Sex Transm Dis* 1994;21:70.

**A randomized, prospective trial comparing amoxicillin and erythromycin for the**

**treatment of *Chlamydia trachomatis* in pregnancy.**

NS SILVERMAN, M SULIVAN, M HOCHMAN, M WOMACK, DL JUNGKIND. *Am J Obstet Gynecol* 1994;170:829.

**Urine as an alternative to urethral swabs for the diagnosis of *Chlamydia trachomatis* in infertile males.**

TN RAMUTHAGA, MS BORNMAN, M MAHOMED, D BOOMKER, AS GREEF, HH CREWEBROWN, S REIF. *Int J Andrology* 1994;17:9.

**Non-invasive sampling for detection of genital infection with *Chlamydia trachomatis* in males utilising urinary leukocyte esterase tests and immunoassays.**

MA DOMEIKA, M BASSIRI, PA MARDH. *Infection* 1994;22:65.

**Oral contraceptives do not alter the course of experimentally induced chlamydial salpingitis in monkeys.**

DL PATTON, YTC SWEENEY, CC KUO. *Sex Transm Dis* 1994;21:89.

**Antibody recognition of a neutralization epitope on the major outer membrane protein of *Chlamydia trachomatis*.**

GM ZHONG, J BERRY, RC BRUNHAM. *Infect Immun* 1994;62:1576.

**Toxic effect on human spermatozoa by *Chlamydia trachomatis* purified lipopolysaccharide.**

F GALDIERO, L SOMMESE, F GORGA, E GALDIERO, A RIZZO, M AJELLO. *Fems Microbiol Lett* 1994;115:197.

**Interaction of the *Chlamydia trachomatis* histone H1-like protein (Hc1) with DNA and RNA causes repression of transcription and translation in vitro.**

LB PEDERSEN, S BIRKLUND, G CHRISTIANSEN. *Mol Microbiol* 1994;11:1085.

### Pelvic inflammatory disease

**Epidemiology of pelvic inflammatory disease.**

F GRODSTEIN, KJ ROTHMAN. *Epidemiol* 1994;5:234.

**Observations concerning the microbial etiology of acute salpingitis.**

DE SOPER, NJ BROCKWELL, HP DALTON, D JOHNSON. *Am J Obstet Gynecol* 1994;170:1008.

### Candidiasis

**Detection of anti-*Candida albicans* IgE antibodies in vaginal washes from patients with acute vulvovaginal candidiasis.**

P REGULEZ, JFG FERNANDEZ, MD MORAGUES, J SCHNEIDER, G QUINDOS, J PONTON. *Gynecol Obstet Invest* 1994;37:110.

**Effects of preinduced candida-specific cell-mediated immunity on experimental vaginal candidiasis.**

PL FIDEL, ME LYNCH, JD SOBEL. *Infect Immun* 1994;62:1032.

**Adherence of *Candida albicans* to endothelial cells is inhibited by prostaglandin I-2.**

SA KLOTZ. *Infect Immun* 1994;62:1497.

**Mechanisms by which *Candida albicans* induces endothelial cell prostaglandin synthesis.**

SG FILLER, BO IBE, AS IBRAHIM, MA GHANNOUM, JV RAJ, JE EDWARDS. *Infect Immun* 1994;62:1064.

**Idiotypic intravaginal vaccination to protect against vaginitis by secretory, yeast killer toxin-like anti-idiotypic antibodies.**

L POLONELLI, F DEBERNARDIS, S CONTI, *et al.* *J Immunol* 1994;152:3175.

**The role of microfilaments and microtubules during pH-regulated morphological transition in *Candida albicans*.**

K YOKOYANA, H KAJI, K NISHIMURA, M MIYAJI. *Microbiology* 1994;140:281.

### Bacterial vaginosis

**Bacterial vaginosis—efficacy and safety of intravaginal metronidazole treatment.**

CH LIVENGOD, JA MCGREGOR, DE SOPER, E NEWTON, JL THOMASON. *Am J Obstet Gynecol* 1994;170:759.

**Bacterial vaginosis is associated with prematurity and vaginal fluid mucinase and sialidase: results of a controlled trial of topical clindamycin cream.**

JA MCGREGOR, JI FRENCH, W JONES, *et al.* *Am J Obstet Gynecol* 1994;170:1048.

### Trichomoniasis

**A new method for identification of *Trichomonas vaginalis* is by fluorescent DNA insitu hybridization.**

R MURESU, S RUBINO, P RIZZU, A BALDINI, M COLOMBO, P CAPPUCINELLI. *J Clin Microbiol* 1994;32:1018.

### Syphilis and other treponematoses

**Epidemic syphilis: maternal factors associated with congenital infection.**

BL MCFARLIN, SF BOTTOMS, BS DOCK, NB ISADA. *Am J Obstet Gynecol* 1994;170:535.

**A case of secondary syphilis combining classical and atypical features.**

Z DAMANHOURY, EL SMITH. *Eur J Dermatol* 1994;4:159.

**HIV and syphilis seroprevalence among clients with sexually transmitted diseases attending a walk-in clinic at Cook County Hospital.**

DA ANSELL, TC HU, M STRAUS, M COHEN, R SHERER. *Sex Transm Dis* 1994;21:93.

**Evaluation of neurosyphilis in human immunodeficiency virus infected individuals.**

MG TOMBERLIN, PD HOLTOM, JL OWENS, RA LARSEN. *Clin Infect Dis* 1994;18:288.

**Diagnosing neurosyphilis—response.**

EW HOOK. *Clin Infect Dis* 1994;18:295.

**Immunisation of guinea pigs with *Treponema pallidum* recombinant antigens reveals the presence of novel epitopes.**

K WICHER, F ABBRUSCATO, V WICHER, L SCHOULS. *Int Arch Allergy Immunol* 1994;103:396.

## Hepatitis

### Research in chronic viral hepatitis.

*Archives of Virology Supplement* 8; 1993 (whole issue).

### Spousal transmission of the hepatitis C virus?

LB SEEFF, HJ ALTER. *Ann Intern Med* 1994;120:807.

### Sexual transmission of hepatitis C.

SJ SKIDMORE, KE COLLINGHAM, SM DRAKE. *J Med Virol* 1994;42:247.

### Hepatitis C virus infection in a sexually active inner city population. The potential for heterosexual transmission.

GL DAIKOS, S LAI, MA FISCHI. *Infection* 1994;22:72.

### Hepatitis B virus infection in a group of heterosexuals with multiple partners in Amsterdam, The Netherlands: implications for vaccination?

GJJ VANDOORNUM, HJA VANHAASSTRECHT, C HODYKASS, et al. *J Med Virol* 1994;43:20.

## Herpes

### Analysis of the epidemiology and pathogenesis of herpes simplex virus infections in pregnant women and infants using the HSV-2 glycoprotein G antibody assay.

AM ARVIN, CG PROBER. *Infect Agents Dis* 1994;2:375.

### Seroprevalence of antibodies to human herpes viruses in England and Hong Kong.

*J Med Virol* 1994;43:91.

### Autopsy findings in two cases of neonatal herpes simplex virus infection: detection of virus by immunohistochemistry, in situ hybridization and the polymerase chain reaction.

JAR NICOLL, S LOVE, PA BURTON, PJ BERRY. *Histopathol* 1994;24:257.

### Diagnosis of genital herpes by polymerase chain reaction amplification.

T GOTO, Y YAMAGUCHI, M HASHIDO, H YOSHIKAWA, T KAWANA. *Microbiol Immunol* 1994;37:987.

### Acyclovir. Is the honeymoon coming to an end?

PD WELSBY. *J Infect* 1994;28:121.

### Genital herpes—a guide to pharmacological therapy.

A DE RUITER, RN THIN. *Drugs* 1994;47:297.

### Novel carbohydrate conjugates as potential prodrugs of acyclovir.

SD CHAMBERLAIN, AR MOORMAN, TC BURNETTE, P DEMIRANDA, TA KRENITSKY. *Antivir Chem Chemother* 1994;5:64.

### Antigenic specificities of human CD4 (+) T-cell clones recovered from recurrent genital herpes simplex virus type 2 lesions.

DM KOELLE, L COREY, RL BURKE, et al. *J Virol* 1994;68:2803.

### Mannan-binding protein and bovine conglutinin mediate enhancement of herpes simplex virus type 2 infection in mice.

PB FISCHER, S ELLERMANNRIKEN, S THIEL, JC JENSENTUS, SC MOGENSEN. *Scand J Immunol* 1994;39:439.

### The transmembrane domain of the large subunit of HSV-2 ribonucleotide reductase (ICP10) is required for protein kinase activity and transformation-related signaling pathways that result in ras activation.

CC SMITH, JH LUO, JCR HUNTER, JV ORDONEZ, L AURELIAN. *Virology* 1994;200:598.

## Human papillomavirus infection

### The absence of vaginal human papillomavirus 16 DNA in women who have not experienced sexual intercourse.

E RYLANDER, L RUUSUVAARA, MW ALMSTROMER, M EVANDER, G WADELL. *Obstet Gynecol* 1994;83:735.

### Low concordance of genital human papillomavirus lesions and viral types in HPV-infected women and their male sexual partners.

MI HIPPELAINEINEN, M YLISKOSKI, S SYRJANEN, et al. *Sex Transm Dis* 1994;21:76.

### Detection of human papillomavirus by the polymerase chain reaction in histologically normal penile skin adjacent to penile warts.

KA WARD, PC WINTER, M WALSH, RD MAW, WW DINSMORE. *Sex Transm Dis* 1994;21:83.

### Urethral meatal warts in men; results of urethroscopy and biopsy.

I ROTHMAN, RE BERGER, N KIVIAT, AL NAVARRO, ML REMINGTON. *J Urol* 1994;151:875.

### Treatment of genital HPV infection with carbon dioxide laser and systemic interferon alpha-2b.

P NIEMINENE, M AHO, M LEHTINEN, E EVESTERINEN, A VAHERI, J PAAVONEN. *Sex Transm Dis* 1994;21:65.

### Prevalence of anal human papillomavirus infection and intraepithelial neoplasia in renal allograft recipients.

OA OGUNBIYI, JH SCHOLEFIELD, AT RAFTERY, et al. *Br J Surg* 1994;81:365.

### Risk of genital human papillomavirus infection in women with human immunodeficiency virus induced immunosuppression.

GYF HO, RD BURK, I FLEMING, RS KLEIN. *Int J Cancer* 1994;56:788.

### Typing of human papillomaviruses in condylomata acuminata from Greece.

V LABROPOULOU, A BALAMOTIS, A TOSCA, A ROTOLA, P MAVROMARANAZOS. *J Med Virol* 1994;42:259.

### Detection of human papillomavirus L1 protein in condylomata acuminata from various anatomical sites.

K WOOLS, JT BRYAN, BP KATZ, M RODRIQUEZ, T DAVIS, DR BROWN. *Sex Transm Dis* 1994;21:103.

### Prevalence of HPV in premalignant and malignant cervical lesions in Greenland and Denmark; PCR and insitu hybridization analysis on archival material.

AM SEBBELOV, C SVENDSEN, H JENSEN, SK KJAER, B NORRILD. *Res Virol* 1994;145:83.

### Human papillomavirus infection in Norwegian women with cervical cancer.

E KRISTIANSEN, A JENKINS, G KRISTENSEN, et al. *APMIS* 1994;102:122.

### Human papillomavirus and cervical cancer in Honduran women.

W MELCHERS, A FERRERA, D WILLEMSE, et al. *Am J Trop Med Hygiene* 1994;50:137.

### Multivariate discriminant analysis of normal, intraepithelial neoplasia and human papillomavirus infection of the uterine cervix samples.

E ARTACHOPERULO, R ROLDANCILLALOBOS, J SALASMOLINA, R VAAMONDELEMO. *Histol Histopathol* 1994;9:135.

### Sexually transmitted diseases and other risk factors for cervical dysplasia among southwestern Hispanic and non-Hispanic white women.

TM BECKER, CM WHEELER, NS MCGOUGH, et al. *JAMA* 1994;271:1181.

### Human papillomavirus in non-genital Bowen's disease and Bowenoid carcinoma.

A VENUTI, P DONATI, A AMANTEA, L BALUS. *Eur J Dermatol* 1994;4:142.

### Absence of human papillomavirus in squamous cell carcinomas of nongenital skin from immunocompromised renal transplant patients.

SE SMITH, IC DAVIS, B LESHIN, AB FLEISCHER, WL WHITE, SR FELDMAN. *Arch Dermatol* 1994;129:1585.

### Analysis of cytomorphologically abnormal cervical scrapes for the presence of 27 microtropic human papillomavirus genotypes, using polymerase chain reaction.

AMD HUSMAN, JMM WALBOOMERS, CJLM MEIJER, et al. *Int J Cancer* 1994;56:802.

### Detection of human papillomaviruses in paraffin-embedded biopsies of cervical intraepithelial lesions; analysis by immunohistochemistry, in situ hybridization and the polymerase chain reaction.

P DELVENNE, MA FONTAINE, C DELVENNE, A NIKKELS, J BONIVER. *Modern Pathol* 1994;7:113.

### Human papillomavirus and epithelial carcinogenesis.

S EUVRARD, J VIAC, Y CHARDONNET. *Eur J Dermatol* 1994;4:138.

### Genital papillomaviruses and related neoplasms; causation, diagnosis and classification (Bethesda).

CP CRUM. *Modern Pathol* 1994;7:138.

### Disrupted dichotomous intracellular control of human papillomavirus infection in cancer of the cervix.

H ZUR HAUSEN. *Lancet* 1994;343:955.

### p53 mutations in HPV-negative cervical carcinoma.

DJ PARK, SP WILCZYNSKI, RL PAQUETTE, CW MILLER, HP KOEFFLER. *Oncogene* 1994;9:205.

### p53 mutations in cervical carcinogenesis low frequency and lack of correlation with human papillomavirus status.

RMC BUSBYEARLE, CM STEEL, ARW WILLIAMS, B COHEN, CC BIRD. *Br J Cancer* 1994;69:732.

### Coexistence of episomal and integrated HPV16 DNA in squamous cell carcinoma of the cervix.

E KRISTIANSEN, A JENKINS, R HOLM. *J Clin Pathol* 1994;47:253.

### Genome amplification of human papillomavirus types 16 and 18 in cervical carcinomas is related to the retention of E1/E2 genes.

J BERUMEN, L CASAS, E SEGURA, JL AMEZCUA, A GARCIAARRANCA. *Int J Cancer* 1994;56:640.

### Cervical dysplasia regression induced by all-trans-retinoic acid.

MB SPORN, AB ROBERTS. *J Nat Cancer Inst* 1994;86:476.

**Enhancement of regression of cervical intraepithelial neoplasia-II (moderate dysplasia) with topically applied all-trans-retinoic acid—a randomized trial.**  
FL MEYSKENS, E SURWIT, TE MOON, *et al.* *J Nat Cancer Inst* 1994;86:539.

**A virus-like particle enzyme-linked immunosorbant assay detects serum antibodies in a majority of women infected with human papillomavirus type-16.**  
R KIRNBAUER, NL HUBBERT, CM WHEELER, TM BECKER, DR OWYM, JT SCHILLER. *J Nat Cancer Inst* 1994;86:494.

**Papillomavirus capsids—a new approach to identify serological markers of HPV infection.**  
DA GALLOWAY. *J Nat Cancer Inst* 1994;86:474.

**Retention of low copy number human papillomavirus DNA in cultured cutaneous and mucosal wart keratinocytes.**  
AT WILLIAMS, CJ SEXTON, KJ PURDIE, *et al.* *J Gen Virol* 1994;75:505.

**Interferon and retinoic acid suppress the growth of human papillomavirus type 16 immortalized cervical epithelial cells, but only interferon suppresses the level of human papillomavirus transforming oncogenes.**  
C AGARWAL, JR HEMBREE, EA RORKE, RL ECKERT. *Cancer Res* 1994;54:2108.

**Adeno-associated virus inhibits human papillomavirus type 16; a viral interaction implicated in cervical cancer.**  
PL HERMONAT. *Cancer Res* 1994;54:2278.

**Interaction of the E6 protein of human papillomavirus with cellular proteins.**  
N KEEN, R ELSTON, L CRAWFORD. *Oncogene* 1994;9:1493.

**Sequence rearrangements in the upstream regulatory region of human papillomavirus type 6; are these involved in malignant transition.**  
H KITSATO, H DELUS, H ZURHAUSEN, K SORGER, F ROSL, EM DEVILLIERS. *J Gen Virol* 1994;75:1157.

**Papillomavirus containing cis-acting sequences that can suppress but not regulate origins of DNA replication.**  
FS NALLASETH, ML DEPAMPHILIS. *J Virol* 1994;68:3051.

**The E6/E7 promoter of extrachromosomal HPV16 DNA in cervical cancers escapes from cellular repression by mutation of target sequences for YY1.**  
M MAY, XP DONG, E BEYERFINKLER, F STUBEN-RAUCH, PG FUCHS, H PFISTER. *EMBO J* 1994;13:1460.

**Regulation of early gene expression of human papillomavirus type 16 by inflammatory cytokines.**  
S KYO, M INOUE, N HAYASAKA, T INOUE, M YUTSUDO, O TANIZAWA, A HAKURA. *Virology* 1994;200:130.

**Sequence variation in the capsid protein genes of human papillomavirus type 16.**  
P PUSHKO, T SASAGAWA, J CUZICK, J CRAWFORD. *J Gen Virol* 1994;75:911.

**Reversible repression of papillomavirus oncogene expression in cervical carcinoma cells; consequences for the phenotype and E6-p53 and E7-pRB interactions.**  
M VONKNEBELDOEBERTZ, C RITTMULLER, F AENGENEYNDT, P JANDENDURR, D SPITKOVSKY. *J Virol* 1994;68:2811.

**Differentiation-independent constitutive expression of the human papillomavirus type 16 E6 and E7 oncogenes in the CaSki cervical tumour cell line.**

CK CHOO, EA RORKE, RL ECKERT. *J Gen Virol* 1994;75:1139.

**“Natural” presentation of human papillomavirus type-16 E7 protein to immunocompetent mice results in antigen-specific sensitization or sustained unresponsiveness.**  
MA CHAMBERS, Z WEI, N COLEMAN, AA NASH, MA STANLEY. *Eur J Immunol* 1994;24:738.

**Correlation between tumorigenicity and expression levels or splicing patterns of transcripts of the human papillomavirus type 16 E6 gene.**  
T INOUE, S KYO, T KIYONO, *et al.* *Jpn J Cancer Res* 1994;85:357.

**Synergistic effect of retinoids and interferon  $\alpha$  on tumor-induced angiogenesis; anti-angiogenic defect on HPV-harboring tumor-cell lines.**  
D ROSSI, S FUMAGALLI, L VOLPI, L LARIZZA. *Int J Cancer* 1994;57:81.

**Self-assembly of human papillomavirus type 16 capsids by expression of the L1 protein in insect cells.**  
P LECANN, P COURSAGET, S IOCHMANN, A TOUZE. *Fems Microbiol Lett* 1994;117:269.

## Cervical cytology and colposcopy

**Improving accuracy in gynecologic cytology—results of the College of American Pathologists interlaboratory comparison program in cervicovaginal cytology.**  
DD DAVEY, MML NIELSEN, WJ FRABLE, W ROSENSTOCK, SM LOWELL, BB KRAEMER. *Arch Pathol Lab Med* 1994;117:1193.

**Cytological changes preceding cervical cancer.**  
JH ROBERTSON, B WOODEND, H ELLIOTT. *J Clin Pathol* 1994;47:278.

**Sensitivity of the Papanicolaou smear in human immunodeficiency virus infected women.**  
AP KORN, M AUTRY, PA DEREMER, W TAN. *Obstet Gynecol* 1994;83:401.

**Cervicovaginal screening in women with HIV infection; a need for increased vigilance?**  
CA HANKINS, JA LAMONT, MA HANDLEY. *Can Med Assoc J* 1994;150:681.

**Cytologic detection of *Strongyloides stercoralis* in a routine cervicovaginal smear.**  
DA MURTY, UK LUTHRA, K SEHGAL, P SODHANI. *Acta Cytol* 1994;38:223.

**Thin Prep processor—clinical trials demonstrate an increased detection rate of abnormal cervical cytologic specimens.**  
DC WILBUR, ES CIBAS, S MERRITT, LP JAMES, BM BERGER, TA BONFIGLIO. *Am J Clin Pathol* 1994;101:209.

**Homogenous sampling accounts for the increased diagnostic accuracy using the Thin Prep processor.**  
ML HITCHINSON, LM ISENSTEIN, A GOODMAN, *et al.* *Am J Clin Pathol* 1994;101:215.

**Evaluation of the PAPNET cytologic screening system for quality control of cervical smears.**  
LG KOSS, E LIN, K SCHREIBER, P ELGERT, L MANGO. *Am J Clin Pathol* 1994;101:220.

**Adolescents and colposcopy; The use of different procedures to reduce anxiety.**

VI RICKERT, KJ KOZLOWSKI, AM WARREN, A HENDON, P DAVIS. *Am J Obstet Gynecol* 1994;170:504.

**Digital imaging colposcopy, image analysis and quantification of the colposcopic image.**  
MI SHAFI, JA DUNN, EJ BUXTON, C WILLIAMS, DM LUESLEY. *Br J Obstet Gynaecol* 1994;101:234.

## Other sexually transmitted diseases

**Proceedings of the 10th International Meeting, International Society for Sexually Transmitted Diseases Research, Helsinki, Finland 1993.**  
*Sex Trans Dis* 1994;21 Suppl Mar-Apr (whole issue)

**Gross and microscopic features in chancroid; a study in 200 new culture proven cases in San Salvador.**  
C ORTIZZEPEDA, E HERNANDEZPEREZ, R MARROQUINBURGOS. *Sex Transm Dis* 1994;21:112.

## Public health and social aspects

**A comparison of risky health behaviours of sexually active, sexually abused and abstaining adolescents.**  
S NAGY, AG ADCOCK, MC NAGY. *Pediatrics* 1994;93:570.

**Does a condom-promoting strategy (the Swiss-stop-AIDS-campaign) modify sexual behaviour among adolescents.**  
D HAUSSER, PA MICHAUD. *Pediatrics* 1994;93:580.

## Microbiology and Immunology

**Secretory immunity in the female reproductive tract.**  
WH KUTTEH, J MESTECKY. *Am J Reprod Immunol* 1994;31:40.

**Mixed-effect models for predicting microbial interactions in the vaginal ecosystem.**  
RA ROSS, MLT LEE, ML DELANEY, AB ONDERDONK. *J Clin Microbiol* 1994;32:871.

**Vaginal lactobacillosis.**  
BJ HOROWITZ, PA MARDH, E NAGY, EL RANK. *Am J Obstet Gynecol* 1994;170:857.

**New strategies for using mucosal vaccination to achieve more effective immunisation.**  
RI WALKER. *Vaccine* 1994;12:387.

**Identification of five peptostreptococcus species isolated predominantly from the female genital tract by using the rapid ID32A system.**  
J NG, LK NG, AW CHOW, JAR DILLON. *J Clin Microbiol* 1994;32:1302.

## Dermatology

**Lichen sclerosus—a review.**  
CM RIDLEY. *Eur J Dermatol* 1994;4:99.

**Surgical treatment of idiopathic calcinosis of the scrotum.**

FWB PREIS, M CAVALLINI, A TODDE, M COLONNA, E SERVIDA. *Eur J Dermatol* 1994;4:226.

**Penile syringoma: a case report.**

RC RATNAVEL, RN LONSDALE, PG NORRIS. *Eur J Dermatol* 1994;4:229.

---

## Miscellaneous

---

**The incidence of genitourinary infection in a cohort of healthy women.**

WM McCORMACK, SH ZINNER, WM McCORMACK. *Sex Transm Dis* 1994;21:63.

**Diagnostic tests for sexually transmitted diseases; a challenge.**

S BERKLEY. *Lancet* 1994;343:685.

**Sexually transmitted diseases in the tropics, 30.**

PL PERINE. *Med J Australia* 1994;160:358.

**Treatment regimens of STD patients in Cameroon; a need for intervention.**

A TREBUCQ, JP LOUIS, JP TCHUPO, R MIGLIANI, J SMITH, E DELAPORTE. *Sex Transm Dis* 1994;21:124.

**Circumcision and sexually transmitted diseases.**

LS COOK, LA KOUTSKY, KK HOLMES. *Am J Publ Health* 1994;84:197.

**Reactive arthritis.**

B SVENUNGSSON. *BMJ* 1994;308:671.

**Challenges for the development of female-controlled vaginal microbicides.**

CJ ELIAS, LL HEISE. *AIDS* 1994;8:1.

**Urinary tract infection in men—state of the art.**

AJ SCHAEFFER. *Infection* 1994;22:S19.

**On the relevance of Gram-positive bacteria in prostatitis.**

B BERGMAN. *Infection* 1994;22:S22.

**Complicated UTI and prostatitis in men.**

A PFAU. *Infection* 1994;22:S58.

**Carcinoma in situ of the penis.**

GS GERBER. *J Urol* 1994;151:829.

**Carcinoma of the penis—editorial.**

WS McDOUGAL. *J Urol* 1994;151:1250.

**Lymphoma of genitourinary tract.**

YH SALEM, HC MILLER. *J Urol* 1994;151:1162.

**Genital examination under ketamine sedation in cases of suspected sexual abuse—commentary.**

CJ HOBBS. *Arch Dis Child* 1994;70:197.

**Repeatability of interview derived information on sexual history: a study in women.**

TE ROHNA, JR MCLAUGHLIN, DG HARNISH. *Epidemiology* 1994;5:360.